



Neuromodulator Post Care

- Visible bumps may be seen at the injection sites. These are normal and may last up to a few hours.
- The product will start to work in 2-7 days and peak in about 2 weeks. Please note, if additional product is needed, it will not be injected until about 14 days after your initial injection.
- Do not lie down for 4 hours post injection.
- Do not massage the treatment area.
- Do not do strenuous exercise for 24-48 hours following your injection.
- Do not get a massage, do microdermabrasion or any other activity that may elevate your core body temperature.
- If forehead was treated, do not wear a hat or headband for 48 hours.
- You may cleanse your face normally this evening but please do not massage the injected area. Also, do not use any abrasive exfoliants or scrubs.
- Bruising may be seen. This happens in a small percentage of treatments but will not affect your treatment results. There also may be some discomfort associated with it. Bruising should resolve within one week.
- Cosmetics, moisturizers, lotions may be used the next day, but do not apply extreme pressure or do anything that may cause discomfort.
- Exercising the muscles in the treated area may help the product get into the muscle faster, but has no effect on the overall efficacy of the treatment.
- Headache may be a possible side effect. Take Motrin or Tylenol for your headache should it occur.
- Sensitive or blurry vision is possible for 2-3 days post treatment. Wear sunglasses and moisturize eyes with eye drops if needed.
- Flu-like symptoms are rare, but may last 2-3 days. Fluids, rest, and Motrin/Tylenol may be used.
- To protect your skin and reduce the effects of aging, we recommend sunscreen SPF 45 or greater be applied to exposed skin every 4 hours during sun exposure.
- **Any symptoms or problems other than those listed should be reported immediately to Lift Aesthetics, LLC: 503-810-7281**

Touch-Up Policy for Neuromodulator:

- After two weeks mild asymmetry may be noticeable.
- Do not panic, this is normal and can usually be corrected with a touch-up treatment.